

Proof

Max cardinality matching in $G \le \text{value of maxflow in } G'$

- ightharpoonup Given a maximum matching M of cardinality k.
- ightharpoonup Consider flow f that sends one unit along each of k paths.
- \blacktriangleright f is a flow and has cardinality k.





© Harald Räcke

13.1 Matching

477

479

13.1 Matching


Which flow algorithm to use?


- ▶ Generic augmenting path: $\mathcal{O}(m \operatorname{val}(f^*)) = \mathcal{O}(mn)$.
- ► Capacity scaling: $\mathcal{O}(m^2 \log C) = \mathcal{O}(m^2)$.

Proof

Max cardinality matching in $G \ge \text{value of maxflow in } G'$

- Let f be a maxflow in G' of value k
- ▶ Integrality theorem $\Rightarrow k$ integral; we can assume f is 0/1.
- ▶ Consider M= set of edges from L to R with f(e) = 1.
- ► Each node in *L* and *R* participates in at most one edge in *M*.
- ▶ |M| = k, as the flow must use at least k middle edges.





© Harald Räcke

13.1 Matching

478

Baseball Elimination

team	wins	losses	remaining games			
i	w_i	ℓ_i	Atl	Phi	NY	Mon
Atlanta	83	71	_	1	6	1
Philadelphia	80	79	1	_	0	2
New York	78	78	6	0	_	0
Montreal	77	82	1	2	0	_

Which team can end the season with most wins?

- Montreal is eliminated, since even after winning all remaining games there are only 80 wins.
- But also Philadelphia is eliminated. Why?

Baseball Elimination

Formal definition of the problem:

- ▶ Given a set S of teams, and one specific team $z \in S$.
- ▶ Team x has already won w_x games.
- ▶ Team x still has to play team y, r_{xy} times.
- ▶ Does team z still have a chance to finish with the most number of wins.

☐☐☐☐☐☐ © Harald Räcke

13.2 Baseball Elimination

Certificate of Elimination


Let $T \subseteq S$ be a subset of teams. Define

$$w(T) := \sum_{i \in T} w_i, \qquad r(T) := \sum_{i,j \in T, i < j} r_{ij}$$
 wins of teams in T

If $\frac{w(T)+r(T)}{|T|} > M$ then one of the teams in T will have more than M wins in the end. A team that can win at most M games is therefore eliminated.

Baseball Elimination

Flow network for z = 3. M is number of wins Team 3 can still obtain.



Idea. Distribute the results of remaining games in such a way that no team gets too many wins.

Marald Räcke

13.2 Baseball Elimination

Theorem 1

A team z is eliminated if and only if the flow network for z does not allow a flow of value $\sum_{i,j \in S \setminus \{z\}, i < j} \gamma_{i,j}$.

Proof (⇐)

- ► Consider the mincut A in the flow network. Let T be the set of team-nodes in A.
- If for node x-y not both team-nodes x and y are in T, then $x-y \notin A$ as otw. the cut would cut an infinite capacity edge.
- ▶ We don't find a flow that saturates all source edges:

$$r(S \setminus \{z\}) > \operatorname{cap}(A, V \setminus A)$$

$$\geq \sum_{i < j: i \notin T \lor j \notin T} r_{ij} + \sum_{i \in T} (M - w_i)$$

$$\geq r(S \setminus \{z\}) - r(T) + |T|M - w(T)$$

▶ This gives M < (w(T) + r(T))/|T|, i.e., z is eliminated.

Baseball Elimination

Proof (⇒)

- ▶ Suppose we have a flow that saturates all source edges.
- ▶ We can assume that this flow is integral.
- ightharpoonup For every pairing x-y it defines how many games team xand team γ should win.
- ightharpoonup The flow leaving the team-node x can be interpreted as the additional number of wins that team x will obtain.
- ▶ This is less than $M w_x$ because of capacity constraints.
- ▶ Hence, we found a set of results for the remaining games, such that no team obtains more than *M* wins in total.
- ▶ Hence, team z is not eliminated.

© Harald Räcke


13.2 Baseball Elimination


485

Project Selection

The prerequisite graph:

- \rightarrow {x, a, z} is a feasible subset.
- \blacktriangleright {x, a} is infeasible.





Project Selection

Project selection problem:

- ▶ Set *P* of possible projects. Project *v* has an associated profit p_{ν} (can be positive or negative).
- Some projects have requirements (taking course EA2 requires course EA1).
- ▶ Dependencies are modelled in a graph. Edge (u, v) means "can't do project u without also doing project v."
- ► A subset A of projects is feasible if the prerequisites of every project in A also belong to A.

Goal: Find a feasible set of projects that maximizes the profit.




13.3 Project Selection

Project Selection

Mincut formulation:

- Edges in the prerequisite graph get infinite capacity.
- Add edge (s, v) with capacity p_v for nodes v with positive profit.
- Create edge (v,t) with capacity $-p_v$ for nodes v with negative profit.



487